

Review

Ferulic acid: pharmaceutical functions, preparation and applications in foods

Shiyi Ou¹ and Kin-Chor Kwok^{2*}

¹Department of Food Science and Technology, Jinan University, Guangzhou, China

²Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong, China

Abstract: Ferulic acid (4-hydroxy-3-methoxycinnamic acid), an effective component of Chinese medicine herbs such as *Angelica sinensis*, *Cimicifuga heracleifolia* and *Lignosticum chuangxiang*, is a ubiquitous phenolic acid in the plant kingdom. It is mainly conjugated with mono- and oligosaccharides, polyamines, lipids and polysaccharides and seldom occurs in a free state in plants. Ferulic acid is a phenolic acid of low toxicity; it can be absorbed and easily metabolized in the human body. Ferulic acid has been reported to have many physiological functions, including antioxidant, antimicrobial, anti-inflammatory, anti-thrombosis, and anti-cancer activities. It also protects against coronary disease, lowers cholesterol and increases sperm viability. Because of these properties and its low toxicity, ferulic acid is now widely used in the food and cosmetic industries. It is used as the raw material for the production of vanillin and preservatives, as a cross-linking agent for the preparation of food gels and edible films, and as an ingredient in sports foods and skin protection agents. Ferulic acid can be prepared by chemical synthesis and through biological transformation. As polysaccharide ferulate is a natural and abundant source of ferulic acid, preparation of ferulic acid from plant cell wall materials will be a prospective pathway.

© 2004 Society of Chemical Industry

Keywords: review; ferulic acid; pharmaceutical functions; metabolism; preparation; applications; antioxidant; antimicrobial; anticancer; preservative; cross-linking agent

INTRODUCTION

Ferulic acid (4-hydroxy-3-methoxycinnamic acid) has received much attention in the study of Chinese medicine since it was found to be one of the effective components in Chinese medicine herbs such as *Angelica sinensis*, *Cimicifuga heracleifolia* and *Lignosticum chuangxiang*.¹ It is a phenolic acid ubiquitously existing in the plant kingdom, which can be absorbed by the small intestine and excreted through the urine.² It is one of the most abundant phenolic acids in plants, varying from 5 g kg⁻¹ in wheat bran to 9 g kg⁻¹ in sugar-beet pulp and 50 g kg⁻¹ in corn kernel.^{3,4} In plants, ferulic acid is rarely found in the free form. It is usually found as ester cross-links with polysaccharides in the cell wall, such as arabinoxylans in grasses, pectin in spinach and sugar beet and xyloglucans in bamboo.⁵ It also can cross-link with proteins.⁶ The cross-linking property of ferulic acid with both polysaccharides and proteins suggests that it can be used in the preparation of complex gels in food applications.

In recent years, there have been an increasing number of reports on the physiological functions of ferulic acid and its derivatives in human. Many applications of ferulic acid in the food industry have also been discovered. This article reviews the physiological role of ferulic acid, its metabolism in humans, its preparation and applications in the food industry.

PHARMACEUTICAL FUNCTIONS

Antioxidant activity

The human body is susceptible to the attack of free radicals and reactive oxygen species leading to a series of diseases, such as atherosclerosis, cancer, cataracts, macula lutea, etc.⁷ The generation of free radicals in the human body is influenced by endogenous and exogenous factors. Superoxide and hydroxyl radicals, which are associated with many diseases, are formed by endogenous processes. Superoxide can be produced by oxidation of the molecules of adrenaline,

*Correspondence to: Kin-Chor Kwok, Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong, China
E-mail: bckckwok@polyu.edu.hk

Contract/grant sponsor: Hong Kong Polytechnic University Research Grant; contract/grant number: A-PC73
(Received 11 November 2003; revised version received 20 February 2004; accepted 20 February 2004)

flavine nucleotides, thiol compounds and sugars in the presence of oxygen.⁸ Exogenous factors such as ultraviolet light, air pollutants and cigarette smoke may also induce the production of free radicals that cause damage to our body. Since free radicals are constantly produced in the human body, a defensive system that can prevent the damage caused by free radicals is essential to human health.

Although endogenous antioxidant systems already exist in the human body, the intake of antioxidants is necessary to strengthen the defensive systems. In recent years, there has been an increasing awareness of the bioactivity and potential health benefits of hydroxycinnamates (eg ferulic, caffeic, *p*-coumaric and sinapic acid) and their conjugates. Free ferulic acid is a good antioxidant since it forms a resonance-stabilized phenoxyl radical. Ferulic acid showed high scavenging activity for hydrogen peroxide, superoxide, hydroxyl radical and nitrogen dioxide free radicals. Ou *et al*⁹ observed that ferulic acid, at a concentration of 250 mg l⁻¹, scavenged 92.5% of hydroxyl radicals in a Fenton reaction system. The nitrogen dioxide radical is known to be toxic agent produced in the metabolism of nitrates and nitrites. Zhang *et al*,¹⁰ using the pulse radiolysis technique, showed that hydroxycinnamic acid derivatives, including ferulic acid, sinapic acid and caffeic acid, scavenged nitrogen dioxide radicals. Ferulic acid can prevent peroxynitrite-mediated nitration of protein-bound and free tyrosine and peroxynitrite-mediated oxidation of dihydrorhodamine 123 and DNA.¹¹ Ferulic acid not only scavenges free radicals, but also increases the activity of enzymes that are responsible for scavenging free radicals and inhibits enzymes that catalyze the production of free radicals. Kayahara *et al*¹² observed that ferulic acid, as well as its amino acid derivatives, showed strong tyrosinase-inhibitory activity and superoxide dismutase-like activity. Kawabata *et al*¹³ found that feeding F334 rats with a dose of 0.1 g kg⁻¹ significantly elevated the activities of detoxifying enzymes, namely glutathione S-transferase in the liver and quinone reductase in the liver and colonic mucosa.

Antioxidant properties of hydroxycinnamates have been measured *in vitro*, and the potential health effects of hydroxycinnamates have been demonstrated in animal models. However, few studies have been undertaken concerning the extent to which these compounds are absorbed *in vivo* from the diet. Further studies are required to show the bioavailability of hydroxycinnamates and to demonstrate that particular ones are responsible for some of the health benefits in humans.

Cholesterol-lowering activity

Ferulic acid and γ -oryzanol have been shown to have cholesterol-lowering activity. γ -Oryzanol is a mixture of ferulic acid esters of sterol and triterpene alcohols. It occurs in rice bran oil at a level of 1–2%, where it serves as a natural antioxidant. Rong *et al*¹⁴ found

that γ -oryzanol treatment resulted in a significant reduction in plasma total cholesterol and non-high-density lipoprotein cholesterol in hamsters fed with diets containing 5% coconut oil and 0.1% cholesterol. In addition, the γ -oryzanol-treated animals also exhibited a reduction in cholesterol absorption versus control animals. Kamal-Eldin *et al*¹⁵ studied the effects of the phenolic compounds butylated hydroxytoluene, sesamin, curcumin and ferulic acid on plasma, liver and lung concentrations of tocopherol and on plasma and liver cholesterol in male Sprague-Dawley rats. Ferulic acid had no effect on the level of tocopherol in the plasma, liver or lung, while the other substances showed different interactions with α - and γ -tocopherols. With the exception of butylated hydroxytoluene, all test compounds tended to decrease total cholesterol in plasma. Ferulic acid tended to lower plasma very-low-density lipoprotein + low-density lipoprotein cholesterol concentrations. It was also found that ferulic acid increased plasma high-density lipoprotein cholesterol while the other compounds reduced it numerically, but not statistically. The results suggest some *in vivo* interactions between the phenolic compounds and tocopherol that may increase the bioavailability of vitamin E and decrease cholesterol in rats. The cholesterol-lowering activity of ferulic acid was confirmed by Kim *et al*.¹⁶ It was suggested that ferulic acid inhibited cholesterol synthesis by competitively inhibiting the activity of hydroxymethylglutaryl CoA reductase in the liver and increasing excretion of acidic sterol.^{16,17}

Prevention against thrombosis and atherosclerosis

Ferulic acid-rich herbs have long been used in China for curing thrombosis. Platelet aggregation, one of the mechanisms involved in repair of blood vessel injury, is related to diseases such as thrombosis. Developing a compound capable of inhibiting platelet aggregation may thus provide a therapeutic means. Kayahara *et al*¹² synthesized three types of ferulic acid derivatives (feruloylaminoacid benzyl or methyl esters, feruloylaminoacids and 4-0-[*N*-(carbobenzyloxy)aminoacyl]ferulic acid). They reported that 4-0-[*N*-(carbobenzyloxy)aminoacyl]ferulic acid (amino acid components: isoleucine, proline) could maintain inhibition of collagen-induced platelet aggregation to the same level as ferulic acid, but showed stronger dissociation of ADP-induced aggregation. In other words, these compounds may not only prevent thrombosis but also dissolve thrombi. From the results of *in vitro* and clinical studies, ferulic acid has been shown to inhibit the activity of thromboxane A₂ (TXA₂) synthetase and blood platelet aggregation.^{18–20} Eicosanoic acid is a precursor of TXA₂. This compound is released from cell membrane lipids by a hydrolysis reaction catalyzed by phospholipase A₂. Pan *et al*²¹ reported that ferulic acid could prevent the release of eicosanoic acid by inhibiting

phospholipase A₂. On the basis of these reports, Ou *et al*¹⁷ summarized that the anti-thrombosis effect of ferulic acid was attributed to its antagonistic effect on TXA₂ and inhibitory effects on TXA₂ synthetase and phospholipase A₂.

As atherosclerosis induced by peroxidation of lipids is a main cause of coronary heart disease, the effects of antioxidants on the development of atherosclerosis in experimental animals have been investigated by some researchers. Hiramatsu *et al*²² tested the antioxidant action of γ -oryzanol in hypercholesterolemic rabbits to determine its effects on the development of atherosclerosis. They found that oleate incorporation into cholesteryl ester by macrophages was significantly reduced in the γ -oryzanol-treated group compared with the non-treated group. Sakamoto *et al*²³ also observed that atherogenic index was reduced by intravenous administration of ferulic acid in a high cholesterol diet fed to rats.

Owing to the antioxidant and cholesterol-lowering activities, as well as the preventive effects against thrombosis and atherosclerosis, ferulic acid is considered a potential chemopreventive agent against coronary heart disease.

Antimicrobial and anti-inflammatory activity

Ferulic acid has been shown to possess inhibitory activity on the growth and reproduction of viruses such as influenza virus, respiratory syncytial virus and AIDS. Ferulic acid and isoferulic acid are the active components of the rhizoma of *Cimicifuga* species, which is used frequently as anti-inflammatory drugs in Japanese Oriental medicines. Hirabayashi *et al*²⁴ investigated the effect of ferulic acid and isoferulic acid on murine interleukin-8 (IL-8) production in response to influenza virus infections *in vitro* and *in vivo* by antibody-sandwich enzyme-linked immunosorbent assay. In the *in vitro* study, they found that both ferulic acid and isoferulic acid reduced the IL-8 levels in the 20-h conditioned medium in comparison with control in a dose-dependent manner. In the *in vivo* study, mice were infected with 1000 PFU (plaque forming units) of the virus and received daily oral administrations of *Cimicifuga heracleifolia* extract (5 mg mouse⁻¹ day⁻¹), ferulic acid (0.5 mg mouse⁻¹ day⁻¹), or isoferulic acid (0.125 mg mouse⁻¹ day⁻¹). The three drugs showed a tendency to reduce IL-8 levels in bronchoalveolar lavage (BAL) obtained 2 days after infection. Moreover, both ferulic acid and isoferulic acid significantly reduced the number of exuded neutrophils into BAL. Sakai *et al*¹ investigated the effect of ferulic acid and isoferulic acid on the production of macrophage inflammatory protein-2 (MIP-2) in a murine macrophage cell line, RAW264.7, in response to respiratory syncytial virus (RSV) infection. In the presence of either ferulic acid or isoferulic acid, RSV-infected cells reduced MIP-2 production in a dose-dependent manner. These results suggest that ferulic acid and isoferulic acid might be responsible, at least in part, for the anti-inflammatory drug effect

of *Cimicifuga heracleifolia* extract through the inhibition of MIP-2 production. Edeas *et al*²⁵ reported that ferulic acid and its derivatives inhibited AIDS virus, suggesting it to be a potential chemopreventive agent for AIDS.

Ferulic acid possesses a wide spectrum of antimicrobial activity. It exhibited antimicrobial activity towards Gram-positive bacteria, Gram-negative bacteria and yeasts.²⁶ It also showed strong inhibitory effects on the growth of some human gastrointestinal microflora, including *Escherichia coli*, *Klebsiella pneumoniae*, *Enterobacter aerogenes*, *Citrobacter koseri*, *Pseudomonas aeruginosa*, *Helicobacter pylori* and *Shigella sonnei*.²⁷⁻²⁹ The antimicrobial mechanism of ferulic acid was attributed to its inhibition of arylamine *N*-acetyltransferase in the bacteria.²⁹ Ferulates of triterpene alcohols and sterols, isolated from the methanol extract of rice bran, had anti-inflammatory activity against inflammation induced by 12-*O*-tetradecanoylphorbol-13-acetate in mice.³⁰

Anti-cancer effect

Studies on the chemopreventive effects of ferulic acid on oral and large bowel carcinogenesis have been reported in recent years. Mori *et al*³¹ studied the effects of ferulic acid on oral cancer in rats by feeding ferulic acid in the diet at a dose of 0.5 g kg⁻¹ after exposure to 4-nitroquinoline-1-oxide (4NQO) for 5 weeks in drinking water at a dose of 0.02 g kg⁻¹. They found that the incidences of tongue carcinomas and preneoplastic lesions (severe dysplasia) were significantly lower on termination of the experiment (32 weeks) than the group with the carcinogen alone. The results suggest that ferulic acid has chemopreventive activity on oral cancer. Recently many researchers have focused their attention on the anti-cancer activity of ferulic acid on colon and rectal cancer.^{13,31,32} Kawabata *et al*¹³ examined the modifying effects of dietary administration of ferulic acid on azoxymethane (AOM)-induced colon carcinogenesis in rats. After 35 weeks, rats which were given ferulic acid during the initiation phase at doses of 0.25 and 0.5 g kg⁻¹, respectively, had lower incidences of colonic carcinomas than those which were given AOM alone. They also observed that ferulic acid could elevate the activities of phase II detoxifying enzymes, glutathione *S*-transferase (GST) and quinone reductase (QR) in liver and colon of rats induced by AOM. These results suggest that detoxifying enzymes are related to the blocking effect of ferulic acid on AOM-induced colon carcinogenesis. As free radicals play an important role in the development of cancer, the anti-cancer effects of some antioxidants is attributed to their ability in scavenging free radicals. Kaul and Khanduja³³ reported that topical application of polyphenols (ellagic acid, tannic acid, caffeic acid and ferulic acid) simultaneously with phorbol-12-myristate-13-acetate or mezerein resulted in significant protection against 7,12-dimethylbenz[a]anthracene-induced skin tumors

in mice. *In vivo* and *in vitro* treatment of murine peritoneal macrophages with the tumor promoters resulted in stimulation of superoxide anion radical formation. In their experiments, the polyphenols were strong inhibitors of the superoxide anion free radical.

Besides the functions mentioned above, ferulic acid has been reported to play important roles in scavenging nitrite,³⁴ as well as increasing sperm viability, trans-membrane migration ratio (TMMR) and the levels of intracellular cAMP and cGMP in spermatozoa.³⁵

Pharmacokinetics of ferulic acid in animals and human

Ferulic acid and other hydroxycinnamic acids occur in plant tissues mainly as low-molecular-weight, water-soluble conjugates present in the cytosol and in bound forms esterified or etherified to cell wall polymers. Water-soluble conjugates are more readily absorbed as ester or the aglycone in the upper digestive tract of non-ruminants. Bound acids are primarily released by microbial action in the hindgut of non-ruminants species, including humans.³⁶

Bourne and Rice-Evans³⁷ investigated the bioavailability of ferulic acid in humans from tomato consumption through the monitoring of the pharmacokinetics of excretion in relation to intake. They found that the peak time for maximal urinary excretion was approximately 7 h and the recovery of ferulic acid in the urine, on the basis of total ferulic acid and feruloyl glucuronide excreted, was 11–25% of the amount ingested. In their subsequent research, no urinary excretion of ferulic acid was detected approximately 10 h after tomato ingestion.³⁸ Chang *et al*³⁹ used pure free ferulic acid to carry out a pharmacokinetic study in Wistar female rats. They concluded that free ferulic acid did not enter enterohepatic circulation. Hence, the chemopreventive effect of ferulic acid on colon carcinogenesis depends largely on the ingestion of fiber-bound ferulic acid in plant tissues, which can get to the colon and is partially released by colon microorganisms.

Absorbed ferulic acid and other flavonoids are present in the rat blood circulation in the form of glucuronide, sulfate and methylate conjugates and are excreted via urine or bile.^{40,41} Piskula *et al*⁴¹ proposed that the first step of conjugation of dietary flavonoids was glucuronidation, occurring in the intestinal mucosa in rats. The glucuronized form entered blood circulation and was sulfated in the liver and methylated in the liver and kidney. The presence of non-conjugated ferulic acid found in rat plasma might be due to overdosing of the animals, as discussed by Piskula and Terao⁴² in their study of quercetin absorption in rats.

From the results mentioned above, it can be concluded that free ferulic acid is absorbed from the alimentary tract and is metabolized into various forms. The range of metabolites and their relative proportions depend on many factors, including dose, route of administration and animal species.

The vast proportion of ferulic acid in plants exists in its bound form in the cell wall cross-linking with polysaccharides and proteins by diferulic acids. The bound form of ferulic acid cannot be absorbed through the gastrointestinal tract wall. Hence, esterase activity is necessary to cleave the ester bonds and to release the free acids into the gut lumen where they then become available for absorption or for further metabolism. It has been demonstrated that microbial esterases present in the intestine of mammals (rats and humans) can release ferulic acid and *p*-coumaric acid into the lumen.^{3,9} Free hydroxycinnamic acids (ferulic, caffeic, *p*-coumaric acids) released from colonic fermentation can be absorbed into the circulatory system.^{3,37,43}

Andreasen *et al*⁴⁴ provided the evidence that diferulic acids could be absorbed via the gastrointestinal tract. The 5-5-, 8-0-4-, and 8-5-diferulic acids were identified in the plasma of rats after oral administration with a mixture of the three acids in oil. Their study also revealed that human and rat colonic microflora containing esterase activity were able to release 5-5-, 8-0-4- and 8-5-diferulic acids from model compounds and dietary cereal brans, hence providing a mechanism for the release of dietary diferulates prior to absorption of the free acids. In addition, cell-free extracts from human and rat small intestine mucosa exhibited esterase activity towards diferulate esters. Hence, esterified diferulates can be released from cereal brans by intestinal enzymes, and those free diferulic acids can be absorbed and enter the circulatory system. However, Kroon *et al*³ showed that, in humans, over 95% of the total release of feruloyl groups took place during fermentation in the colon, and only a small portion was released by gastric and small intestinal treatment. This means that, although the small intestine mucosa secretes esterase, it contributes very little to the release of ferulic acid from the plant cell wall.

In summary, free ferulic acid, its soluble conjugates and bound ones can be absorbed from the human gastrointestinal tract. The first two types can be directly absorbed in the small intestine, but the bound ferulic acid must be first released by microbial esterases in the colon before it can be absorbed. The metabolism, physiological role and pharmacokinetic properties of the released free acid and feruloyl compounds need to be further elucidated in future studies.

PREPARATION OF FERULIC ACID

Preparation of ferulic acid by chemical synthesis

It is well-known that ferulic acid can be prepared by the condensation reaction of vanillin with malonic acid catalyzed by piperidine.⁴⁵ This method produces ferulic acid as a mixture of *trans*- and *cis*-isomers. The yield is high, but it takes as long as three weeks to complete the reaction. Da and Xu⁴⁶ improved this method by using benzylamine as the catalytic agent, methylbenzene as the solvent and a reaction temperature of 85–95 °C. The improved method

increased the yield and reduced the reaction time to 2 h.

Preparation of ferulic acid from natural sources

There are three pathways to prepare ferulic acid from natural resources: (1) from low-molecular-weight ferulic conjugates, (2) from plant cell walls, and (3) by tissue culture or microbial fermentation. Several feruloyl esters of triterpene alcohols and sterols were isolated from the methanol extract of rice bran, of which the main component is γ -oryzanol, accounting for about 1.5–2.8% of rice bran oil by weight.⁴⁷ Taniguchi *et al*⁴⁸ prepared ferulic acid in large quantities from rice bran pitch, a blackish brown waste oil discharged in the process of the production of rice bran oil. In this process, the bran oil waste material was hydrolyzed with sodium hydroxide or potassium hydroxide at 90–100°C for 8 h under atmospheric pressure, producing crude ferulic acid with purity of 70–90%. The solution containing alkaline salt of ferulic acid was acidified with dilute sulfuric acid to precipitate ferulic acid.

There are many reports on the use of feruloyl esterases produced by microorganisms to release ferulic acid from plant cell walls. Feruloyl esterases are a subclass of the carboxylesterases that are able to release ferulic acid from a range of esterified substances including methyl ferulate, feruloylated oligosaccharides and polysaccharides.⁴⁹ These enzymes are secreted by fungal, bacterial and yeast microbes, such as *Aspergillus niger*, *Pycnoporus cinnabarinus*, *Streptomyces avermitilis*, *Clostridium thermocellum*, *Bacillus spp*, *Lactobacilli*, *Pseudomonas fluorescens* and *Breuanomyces anomalus*. Although feruloyl esterases are not yet commercially used to prepare ferulic acids, extensive research studies have been carried out. First, microorganisms that can secrete feruloyl esterases have been screened.^{50–55} Second, some characteristics of the enzymes, such as structure, optimal temperature and pH, factors affecting their stability, and the effect of the substrate and polysaccharide-degrading enzymes on the release of ferulic acid were elucidated.^{52,56,57} Third, the effect of some factors on the production of feruloyl esterases, such as carbon resources, nitrogen resources and ferulic acid were studied.⁵⁸ Fourth, the purification method of ferulic acid from fermentation liquids was established.^{59,60} Future work will be focused on how to improve the production of feruloyl esterases by optimizing fermentation technology and genetic engineering.

Cell suspension cultures of *Beta vulgaris*, *Zea mays* and *Chenopodium rubrum* have been reported to accumulate ferulic acid.^{61–63} Bokern *et al*⁶¹ isolated five water-soluble ferulic conjugates by TLC from callus cultures of *Beta vulgaris*. They were identified as 1-*O*-feruloylglucose, 6-*O*-feruloylsucrose, 6-*O*-hydroxycinnamoylglucosides, 1-*O*-feruloylglucuronosylglycerol and feruloylaspartic acid. These water-soluble conjugates were accumulated in different amount between cell lines and conjugates. The most

frequent one, 1-*O*-feruloylglucose, was accumulated up to 20.0 $\mu\text{mol g}^{-1}$ dry matter. This suggests that screening of cell line can increase the content of ferulic acid conjugates in *Beta vulgaris*.

In contrast to the plant cultures mentioned above, cell cultures of *Ajuga pyramidalis* accumulated free ferulic acid⁶⁴ and also conjugated to anthocyanins⁶⁵ up to 150 mg l^{-1} cell culture on the twelfth day. With the increasing demand for natural ingredients, production of ferulic acid by plant cell culture will become a prospective pathway.

APPLICATIONS IN FOODS

Ferulic acid is of low toxicity with LD₅₀ of 2445 mg kg^{-1} body weight in male and 2113 mg kg^{-1} body weight in female rat.⁶⁶ In Japan, ferulic acid has been approved as a food additive and used as a natural antioxidant in foods, beverages and cosmetics. Also, in the USA and most European countries, numerous medical essences and natural extracts of herbs, coffee, vanilla beans, spices and other botanicals are selected for their high content of ferulic acid and added to foods as an FDA-approved antioxidant concoction.⁶⁷ The main applications are summarized as follows.

Production of vanillin

Vanillin is an important aromatic flavor compound used in foods, beverages, perfume and pharmaceuticals. It is produced on a large scale in the industry through chemical synthesis. However, the vanillin obtained by chemical synthesis cannot be considered as natural. Thus, there are many attempts to produce vanillin based on bioconversion of natural sources such as lignin, phenolic stilbenes, isoeugenol, eugenol, ferulic acid or aromatic amino acids.⁶⁸ Ferulic acid can be used to produce vanillin through biotransformation in microorganisms by three major pathways. The first is decarboxylation of ferulic acid by decarboxylase to produce 4-vinylguaicol and then vanillin. The second is the reduction of ferulic acid to dihydroferulic acid, from which vanillic acid and vanillin are formed. The third is to produce vanillic acid and vanillin by formation of coniferyl alcohol from ferulic acid. These biotransformations have been discovered in many bacteria, fungi and yeasts. Use of enzymes secreted by these microorganisms, rather than direct use of microorganisms, is the practical pathway to produce vanillin from ferulic acid, as vanillin is toxic to most microorganisms and the formed vanillin will be further oxidized to the less toxic vanillic acid.⁶⁸

The cell suspension cultures of some plants, such as *Vanilla planifolia* and *Capsicum frutescens*, can also be used to transform ferulic acid to produce vanillin.^{69–71} The pathways involved in the conversion of ferulic acid to vanillin in plant systems have not been investigated in detail, but a pathway analogous to the β -oxidation of fatty acid metabolism was supposed.⁶⁹

Ferulic acid as a preservative

Ferulic acid can be used to preserve foods because of its antioxidant and antimicrobial activities. Ferulic acid was first used in Japan in 1975 to preserve oranges and to inhibit the autooxidation of linseed oil, lard and soybean oil. Mixtures of ferulic acid and amino acids or ferulic acid and dipeptides, such as glycylglycine or alanylalanine, exerted a synergistic inhibitory effect on the peroxidation of linoleic acid.⁶⁷ Ferulic acid is an effective antioxidant in some food-related system, such as lecithin-liposomes and aqueous emulsions.^{72,73}

Compared with other phenolic substances, ferulic acid has two advantages. First, ferulic acid has strong antioxidant activity. Heinonen *et al*⁷⁴ tested the antioxidant activity of gallic acid, propyl gallate, caffeic acid, malvidin, delphinidin, catechin, epicatechin, rutin, quercetin and ferulic acid in lecithin-liposome oxidation systems at doses of 5 and 10 $\mu\text{mol l}^{-1}$. They found that, on the basis of calculation of percentage inhibition values at the propagation phase of oxidation on day 2 for hydroperoxides and on day 3 for hexanal, ferulic acid was the most efficient in inhibiting lipid and protein oxidation. Another advantage is that ferulic acid is much less affected by pH changes than other phenolic compounds, such as chlorogenic acid, caffeic acid and gallic acid.⁷⁵ This is very important for foods subjected to alkali processing, which is commonly used to recover proteins from cereals and legumes, to induce the formation of fiber-forming meat analogue vegetable protein, and to prepare peeled fruits and vegetables. Furthermore, the glucoside esters of ferulic acid also have strong antioxidant activity. The antioxidant activity of some ferulic acid esters is even greater than that of ferulic acid.⁷⁶ This was confirmed by a recent study, in which Kikuzaki *et al*⁷⁷ found that esterification of ferulic acid resulted in increased antioxidant activity, and the activity was influenced by the chain length of the alcohol moiety. When the inhibitory effects of alkyl ferulates against oxidation of liposome induced by 1,1-diphenyl-2-picrylhydrazyl (DPPH) were tested, hexyl, octyl and 2-ethyl-1-hexyl ferulates were more active than the other alkyl ferulates. The results indicated that the affinity of ferulates with lipid substrates affected their antioxidant activity. Medina *et al*⁷⁸ reported that phenolic compounds showed synergistic properties in reinforcing the antioxidant activity of lactoferrin in lipid systems containing iron. However, ferulic acid also has shortcomings. It has been reported that ferulic acid formed 4-vinylguaiacol by decarboxylation during processing, contributing off-flavors in cooked foods, beer and orange juice.^{79,80}

Another preservative function of ferulic acid comes from its antimicrobial activity. Ferulic acid can inhibit the growth of bacteria, fungi and yeasts. Ferulic acid is an active ingredient of extracts of some plants showing antimicrobial activity. Lattanzio *et al*⁸¹ tested the antifungal activity of 12 kinds of phenolic compounds against five fungi (*Sclerotinia sclerotiorum*, *Fusarium oxysporum*, *Alternaria* sp, *Botrytis*

cinerea and *Penicillium digitatum*) and found that ferulic acid showed the highest activity. It was reported that feruloyl oligosaccharide ester showed inhibitory effects against many kinds of Gram-positive and -negative bacteria.⁸² Ferulic acid, at a concentration of 500 mg l^{-1} , can also appreciably inhibit growth of yeasts such as *Pichia anomala*, *Debaryomyces hansenii* and *Saccharomyces cerevisiae*, but it is less effective than potassium sorbate.⁸³

As a cross-linking agent

As ferulic acid is known to cross-link with polysaccharides, it is used to increase the viscosity and form gels from some polysaccharides such as pectin and arabinoxylans.⁸⁴⁻⁸⁶ Gel formation can also be effected from feruloylated arabinoxylans and proteins using cross-linking agents such as ammonium sulfate,^{6,87} hydrogen peroxide and peroxidase,⁸⁴ and laccase.⁸⁵ This property is very important, as ferulic acid can make it possible to use polysaccharides of lower molecular weight, which have low viscosity and poor gel formation capacity, to make new gels in food processing.⁶

Ferulic acid and its oxide, quinoid ferulic acid, can react with some amino acids in proteins such as tyrosine, lysine, cysteine^{88,89} and cross-link protein molecules. Thus, it can be used as a cross-linking agent to improve the properties of protein-based edible films. Kwok and Ou⁹⁰ found that incorporation of ferulic acid into the film-forming solution made from soy protein isolate (SPI) increased the tensile strength, the percentage of elongation at break, and decreased the water vapor permeability and gas permeability of the SPI film. O'Connell and Fox⁹¹ reported that ferulic acid at 5.5 mmol l^{-1} markedly enhanced the heat stability of milk at 140 °C. The addition of caffeic acid resulted in a reduction of the reactive lysine and sulphhydryl content and inhibited the dissociation of κ -casein-rich protein from the casein micelles in milk on heating. It was postulated that on heating in milk, caffeic acid was thermally oxidized to quinones which then interacted with nucleophilic amino acid residues to inhibit κ -casein dissociation from the casein micelle. These results suggest that caffeic acid and its related phenolic compounds, such as ferulic acid, may find their prospective applications in the processing of milk products.

As an ingredient in sport foods and skin lotions

As ferulic acid is a potent antioxidant and is believed to stimulate hormone secretion in humans, it is widely used as an ergogenic substance in sport foods.^{92,93}

Ultraviolet light is an exogenous factor that may cause skin damage, resulting in both precancerous and cancerous skin lesions and acceleration of skin ageing.⁹⁴ Reactive oxygen species are believed to be largely responsible for some of the deleterious effects of UV light upon skin. Particularly, prolonged skin exposure to UV light results in a severe decrease of its antioxidant content.⁹⁵ Overproduction of nitric

oxide from keratinocytes seems to play a major role in the integrated response leading to erythema production and inflammation processes following UV radiation exposure.⁹⁶ Owing to the high degree of conjugated unsaturation (as a strong UV absorber) and its permeability to the skin,⁹⁷ ferulic acid constitutes an active ingredient in many skin lotions, sunscreens and hair creams designed for photoprotection⁶⁸ and for skin tumor inhibition.⁹⁸

Concluding remarks

Ferulic acid is beneficial to health due to its chemopreventive activities against coronary heart disease, thrombosis, carcinogenesis and mutagenesis. Many researchers have recently focused their attention on the anti-cancer activity of ferulic acid on colon and rectal cancer. As free ferulic acid does not enter enterohepatic circulation, oral or intravenous ferulic acid does not easily reach the colon. Although dietary fiber-bound ferulic acid can get to the colon and is partly released by colon microorganisms, the concentration of the released ferulic acid is too low to act as a chemoprevention agent.⁹ An approach to solve this problem is to study the synthesis of enzyme-resistant derivatives of ferulic acid which can be delivered to the colon safely. In a recent study, Ou *et al*¹⁷ used maize starch and ferulic acid to synthesize starch ferulate. They found that the starch ferulate was only partly hydrolyzed (less than 10%) by diastase and the bound ferulic acid was largely released by colon microorganisms. Some derivatives of ferulic acid showed even higher biological activities than the free form. For these reasons, synthesis of enzyme-resistant and high-biological-activity derivatives of ferulic acid will be a focused area of future studies. Extensive toxicological and biological experiments are needed to test this kind of product.

Ferulic acid is an abundant ingredient naturally present in the plant kingdom. It can be commercially produced from natural sources. Because of the many health benefits as well as the antioxidant and antimicrobial activity, more extensive uses of ferulic acid in food applications can be anticipated in the future.

ACKNOWLEDGEMENTS

This work was supported by University Research Grant A-PC73 from the Hong Kong Polytechnic University.

REFERENCES

- Sakai S, Kawamata H, Kogure T, Mantani N, Terasawa K, Umatake M and Ochiai H, Inhibitory effect of ferulic acid and isoferulic acid on the production of macrophage inflammatory protein-2 in response to respiratory syncytial virus infection in RAW264.7 cells. *Mediators Inflammation* 8:173-175 (1999).
- Choudhury R, Srini SK, Debnam E and Rice-Evans CA, Urine excretion of hydroxycinnamates and flavonoids after oral and intravenous administration. *Free Radic Bio Med* 27:278-286 (1999).
- Kroon PA, Faulds CB, Ryden P, Robertson JA and Williamson G, Release of covalently bound ferulic acid from fiber in the human colon. *J Agric Food Chem* 45:661-667 (1997).
- Rosazza JPN, Huang Z, Dostal L, Volm T and Rousseau B, Review: Biocatalytic transformations of ferulic acid: an abundant aromatic natural product. *J Ind Microbiol* 15:457-471 (1995).
- Iiyama K, Lam TBT and Stone BA, Covalent cross-links in the cell wall. *Plant Physiol* 104:315-320 (1994).
- Figueroa-Espinoza MC, Morel MH, Surget A, Asther M, Moukha S, Sigoillot JC and Rouau X, Attempt to cross-link feruloylated arabinoxylans and proteins with a fungal laccase. *Food Hydrocolloids* 13:65-71 (1999).
- Young IS and Woodside JV, Antioxidants in health and disease. *J Clin Pathol* 54:176-186 (2001).
- Becker LB, Vanden Hoek TL, Shao ZH, Li CQ and Schumacker PT, Generation of superoxide in cardiomyocytes during ischemia before reperfusion. *Am J Physiol Heart Circ Physiol* 277:H2240-H2246 (1999).
- Ou S, Li Y and Gao K, A study on scavenging activity of wheat bran dietary fiber for free radical. *Acta Nutr Sin* 21:191-194 (1999).
- Zhang ZE, Yao SD, Lin WZ, Wang WF, Jin YZ and Lin NY, Mechanism of reaction of nitrogen dioxide radical with hydroxycinnamic acid derivatives: a pulse radiolysis study. *Free Radic Res* 29:13-16 (1998).
- Ketsawatsakul U, Whiteman M and Halliwell B, A reevaluation of the peroxynitrite scavenging activity of some dietary phenolics. *Biochem Biophys Res Commun* 279:692-699 (2000).
- Kayahara H, Miao Z and Fujiwara G, Synthesis and biological activities of ferulic acid derivatives. *Anticancer Res* 19:3763-3768 (1999).
- Kawabata K, Yamamoto T, Hara A, Shimizu M, Yamada Y, Matsunaga K, Tanaka T and Mori H, Modifying effects of ferulic acid on azoxymethane-induced colon carcinogenesis in F344 rats. *Cancer Lett* 157:15-21 (2000).
- Rong N, Ausman LM and Nicolosi RJ, Oryzanol decreases cholesterol absorption and aortic fatty streaks in hamsters. *Lipids* 32:303-309 (1997).
- Kamal-Eldin A, Frank J, Razdan A, Tengblad S, Basu S and Vessby B, Effects of dietary phenolic compounds on tocopherol, cholesterol, and fatty acids in rats. *Lipids* 35:427-435 (2000).
- Kim HK, Jeong TS, Lee MK, Park YB and Choi MS, Lipid-lowering efficacy of hesperetin metabolites in high-cholesterol fed rats. *Clin Chim Acta: Int J Clin Chem* 327:129-37 (2003).
- Ou S, Bao H and Lan Z, Advances on pharmacological study of ferulic acid and its derivatives. *J Chinese Med Mater* 24:220-221 (2001).
- Xu LN, Xu DC and Zhang ZJ, Investigation on the mechanism of sodium ferulate in lowering blood platelet aggregation: effect of sodium ferulate on equilibrium of TXA₂ and PGI₂. *Acta Acad Med Sinicae* 6:414-417 (1984).
- Wang Z, Gao YH and Huang RS, Sodium ferulate is an inhibitor of TXA₂ synthetase. *Zhongguo Yao Li Xue Bao* 9:430-433 (1988).
- Zhang JJ, Cai Z, Liu XM and Yu SR, The effect of sodium ferulate on TXB₂ and lipid peroxide in patients with acute cerebral infarction. *Acta Univ Med Tongji* 23:103-105 (1994).
- Pan HZ, Ma LM and Xu CM, Antioxidant activity of sodium ferulate for red blood cells membrane. *Integ Trad Chinese Western Med* 5:678-690 (1985).
- Hiramatsu K, Tani T, Kimura Y, Izumi S and Nakane PK, Effect of gamma-oryzanol on atheroma formation in hypercholesterolemic rabbits. *Takai J Exp Clin Med* 15:299-305 (1990).
- Sakamoto K, Tabata T, Shirasaki K, Inagaki T and Nakayama S, Effects of gamma-oryzanol and cycloartenol ferulic acid ester on cholesterol diet induced hyperlipidemia in rats. *Jap J Pharmacol* 45:559-565 (1987).
- Hirabayashi T, Ochiai H, Sakai S, Nakajima K and Terasawa K, Inhibitory effect of ferulic acid and isoferulic acid

- on murine interleukin-8 production in response to influenza-virus infections in-vitro and in-vivo. *Planta Med* 61:221-226 (1995).
- 25 Edeas M, Khalfoun Y, Lazizi Y, Vergnes L, Labidalle S, Postaire E and Lindenbaum A, Effect of the liposolubility of free radical scavengers on the production of antigen P24 from a HIV infected monocytic cell line. *C R Seances Soc Biol Fil* 189:367-73 (1995).
 - 26 Jeong YC, Jae HM and Keun HP, Isolation and identification of 3-methoxy-4-hydroxybenzoic acid and 3-methoxy-4-hydroxycinnamic acid from hot water extracts of *Hovenia dulcis* Thumb and confirmation of their antioxidative and antimicrobial activity. *Korean J Food Sci Technol* 32:1403-1408 (2000).
 - 27 Tsou MF, Hung CF, Lu HF, Wu LT, Chang SH, Chang HL, Chen GW and Chung JG, Effects of caffeic acid, chlorogenic acid and ferulic acid on growth and arylamine N-acetyltransferase activity in *Shigella sonnei* (group D). *Microbios* 101:37-46 (2000).
 - 28 Nilsson CL, Fingerprinting of *Helicobacter pylori* strains by matrix-assisted laser desorption ionization mass spectrometric analysis. *Rapid Commun Mass Spectrom* 13:1061-1071 (1999).
 - 29 Lo HH and Chung JG, The effects of plant phenolics, caffeic acid, chlorogenic acid and ferulic acid on arylamine N-acetyltransferase activities in human gastrointestinal microflora. *Anticancer Res* 19:133-139 (1999).
 - 30 Akihisa T, Yasukawa K, Yamaura M., Ukiya M, Kimura Y, Shimizu N and Arai K, Triterpene alcohol and sterol ferulates from rice bran and their anti-inflammatory effects. *J Agric Food Chem* 48:2313-2319 (2000).
 - 31 Mori H, Kawabara K, Yoshimi N, Tanaka T, Murakami T, Okada T and Murai H, Chemopreventive effects of ferulic acid on oral and rice germ on large bowel carcinogenesis. *Anticancer Res* 19:3775-3778 (1999).
 - 32 Wargovich MJ, Jimenez A, McKee K, Steele VE, Velasco M, Woods J, Price R, Gray K and Kelloff GJ, Efficacy of potential chemopreventive agents on rat colon aberrant crypt formation and progression. *Carcinogenesis* 21:1149-1155 (2000).
 - 33 Kaul A and Khanduja KL, Polyphenols inhibit promotional phase of tumorigenesis: relevance of superoxide radicals. *Nutr Cancer—an International Journal* 32:81-85 (1998).
 - 34 Rundlof T, Olsson E, Wiernik A, Back S, Aune M, Johansson L and Wahlberg I, Potential nitrite scavengers as inhibitors of the formation of N-nitrosamines in solution and tobacco matrix systems. *J Agric Food Chem* 48:4381-4388 (2000).
 - 35 Zheng RL and Zhang H, Effects of ferulic acid on fertile and asthenozoospermic infertile human sperm motility, viability, lipid peroxidation, and cyclic nucleotides. *Free Radic Biol Med* 22:581-586 (1997).
 - 36 Chesson A, Provan GJ, Russell WR, Scobbie L, Richardson AJ and Stewart C, Hydroxycinnamic acids in the digestive tract of livestock and humans. *J Sci Food Agric* 79:373-378 (1999).
 - 37 Bourne LC and Rice-Evans C, Bioavailability of ferulic acid. *Biochem Biophys Res Commun* 253:222-227 (1998).
 - 38 Bourne LC and Rice-Evans C, Detecting and measuring bioavailability of phenolics and flavonoids in humans: pharmacokinetics of urinary excretion of dietary ferulic acid. *Methods Enzymol* 29:91-107 (1999).
 - 39 Chang MX, Xu LY, Chen JS and Feng Y, Pharmacokinetic study of ferulic acid in rats. *J Chinese Med Mater* 18:300-304 (1993).
 - 40 Azuma K, Ippoushi K, Nakayama M, Ito H, Higashio H and Terao J, Absorption of chlorogenic acid and caffeic acid in rats after oral administration. *J Agric Food Chem* 48:5496-5500 (2000).
 - 41 Piskula MK and Terao J, Accumulation of (-)-epicatechin metabolites in rat plasma after oral administration and distribution of conjugation enzymes in rat tissues. *J Nutr* 128:1172-1178 (1998).
 - 42 Piskula MK and Terao J, Quercetin's solubility affects its accumulation in rat plasma after oral administration. *J Agric Food Chem* 46:4313-4317 (1998).
 - 43 Virgili F, Pagana G, Bourne L, Rimbach G, Natella F, Rice-Evans C and Packer L, Ferulic acid excretion as a marker of consumption of a French maritime pine (*Pinus maritima*) bark extract. *Free Radic Biol Med* 28:1249-1256 (2000).
 - 44 Andreasen MF, Kroon PA, Williamson G and Garcia-Conesa M, Intestinal release and uptake of phenolic antioxidant diferulic acids. *Free Radic Biol Med* 31:304-314 (2001).
 - 45 Adams R and Bockstahler TE, Preparation and reaction of hydroxycinnamic acids and esters. *J Amer Chem Soc* 74:5346-5348 (1952).
 - 46 Da YF and Xu YP, Synthesis of trans-ferulic acid. *Chinese J Pharm* 28:188-189 (1997).
 - 47 Zhang XN and Xu PY, An improved technology to extract γ -oryzanol from rice bran oil. *Cereal Feed Ind* 9:36-37 (1997).
 - 48 Taniguchi H, Hosoda A, Tsuno T, Maruta Y and Nomura E, Preparation of ferulic acid and its application for the synthesis of cancer chemopreventive agents. *Anticancer Res* 19:3757-3761 (1999).
 - 49 Kroon PA, Garcia-Conesa MT, Fillingham IJ, Hazlewood GP and Williamson G, Release of ferulic acid dehydrodimers from plant cell walls by feruloyl esterases. *J Sci Food Agric* 79:428-434 (1999).
 - 50 Bonnin E, Brunel M, Gouy Y, Lesage-Meessen L, Asther M and Thibault JF, *Aspergillus niger* I-1472 and *Fynoporus cinnabarinus* MUCL39533, selected for the biotransformation of ferulic acid to vanillin, are also able to produce cell wall polysaccharide-degrading enzymes and feruloyl esterases. *Enzyme Microb Technol* 28:70-80 (2001).
 - 51 Garcia BL, Ball AS, Rodriguez J, Perez-Leblic ML, Arias ME and Copa-Patino JL, Induction of ferulic acid esterase and xylanase activities in *Streptomyces avermitilis* UAH30. *FEMS Microbiol Lett* 158:95-99 (1998).
 - 52 Prates JAM, Tarbouriech N, Charnock SJ, Fontes CMGA, Ferreira LMA and Davies GJ, The structure of the feruloyl esterase module of xylanase 10B from *Clostridium thermocellum* provides insights into substrate recognition. *Structure* 9:1183-1190 (2001).
 - 53 Yu PQ, Maenz DD, McKinnon JJ, Racz VJ and Christensen DA, Release of ferulic acid from oat hulls by *Aspergillus* ferulic acid esterase and *Trichoderma* xylanase. *J Agric Food Chem* 50:1625-1630 (2002).
 - 54 Donaghy J, Kelly PF and Mckay AM, Detection of ferulic acid esterase production by *Bacillus* spp and *Lactobacilli*. *Appl Microbiol Biotechnol* 50:257-260 (1998).
 - 55 Bartolome B, Faulds CB, Kroon PA, Waldron K, Gilbert HJ, Hazlewood G and Williamson G, An *Aspergillus niger* esterase (ferulic acid esterase III) and a recombinant *Pseudomonas fluorescens* subsp cellulosa esterase (XylD) release a 5-5'-ferulic dehydrodimer (diferulic acid) from barley and wheat cell walls. *App Environ Microbiol* 63:208-212 (1997).
 - 56 Williamson G and Vallejo J, Chemical and thermal stability of ferulic acid esterase-III from *Aspergillus niger*. *Int J Biol Macromol* 21:163-167 (1997).
 - 57 Saulnier L, Marot C, Elgorriaga M, Bonnin E and Thibault JF, Thermal and enzymatic treatments for the release of free ferulic acid from maize bran. *Carbohydr Polym* 45:269-275 (2001).
 - 58 Faulds CB, deVries RP, Kroon PA, Visser J and Williamson G, Influence of ferulic acid on the production of feruloyl esterases by *Aspergillus niger*. *FEMS Microbiol Lett* 15:7239-244 (1997).
 - 59 Edlin DAN, Narbad A, Gasson MJ, Dickinson JR and Lloyd D, Purification and characterization of hydroxycinnamate decarboxylase from *Brettanomyces anomalus*. *Enzyme Microb Technol* 22:232-239 (1998).
 - 60 Couteau D and Mathaly P, Fixed-bed purification of ferulic acid from sugar-beet pulp using activated carbon: optimization studies. *Bioresource Technol* 64:17-25 (1998).
 - 61 Bokern M, Heuer S, Wray V, Witte L, Macek T, Vanek T and Strack D, Ferulic acid conjugates and betacyanins from cell cultures of *Beta vulgaris*. *Phytochemistry* 30:3261-3265 (1991).

- 62 Grabber JH, Hatfield RD, Ralph J, Ron J and Arneheim N, Cross-linking in cell walls from maize cell suspensions. *Phytochemistry* 40:1077-1082 (1995).
- 63 Strack D, Bokern M, Berlin J and Sieg S, Metabolic activity of hydroxycinnamic acid glucose esters in cell-suspension cultures of *Chenopodium rubrum*. *Z Naturforsch Section C Biosci* 39:902-907 (1984).
- 64 Madhavi DL, Smith MAL, Linas AC and Mitiku G, Accumulation of ferulic acid in cell cultures of *Ajuga pyramidalis* Metallica Crispa. *J Agric Food Chem* 45:1506-1508 (1997).
- 65 Madhavi DL, Juthangkoon S, Lewen K, Berberjimeze MD and Smith MAL, Characterization of anthocyanins from *Ajuga pyramidalis* Metallica crispa Cell cultures. *J Agric Food Chem* 44:1170-1176 (1996).
- 66 Tada Y, Tayama K and Aoki N, Acute oral toxicity of ferulic acid, natural food additive, in rats. *Ann Rep Tokyo Metr Lab PH* 50:311-313 (1999).
- 67 Graf E, Antioxidant potential of ferulic acid. *Free Radic Biol Med* 13:435-448 (1992).
- 68 Priefert H, Rabenhorst J and Steinbuchel A, Biotechnological production of vanillin. *Appl Microbiol Biotechnol* 56:296-314 (2001).
- 69 Rao SR and Ravishankar GA, Vanilla flavour: production by conventional and biotechnological routes. *J Sci Food Agric* 80:289-304 (2000).
- 70 Funk C and Brodelius PE, Phenylpropanoid metabolism in suspension cultures of *Vanilla planifolia* and induction of vanillic acid formation. *Plant Physiol* 99:256-262 (1992).
- 71 Westcott RJ, Cheetham PSJ and Barraclough AJ, Use of organized viable Vanilla plant aerial roots for the production of natural vanillin. *Phytochemistry* 35:135-138 (1994).
- 72 Frankel EN, Huang SW, Kanner J and German JB, Interfacial phenomena in the evaluation of antioxidants—bulk oils vs emulsions. *J Agric Food Chem* 42:1054-1059 (1994).
- 73 Huang SW, Frankel EN, Aeschbach R and German JB, Partition of selected antioxidants in corn oil-water model systems. *J Agric Food Chem* 45:1991-1994 (1997).
- 74 Heinonen M, Rein D, Satue-Gracia MT, Huang SW, German JB and Frankel EN, Effect of protein on the antioxidant activity of phenolic compounds in a lecithin-liposome oxidation system. *J Agric Food Chem* 46:917-922 (1998).
- 75 Friedman M and Jurgens HS, Effect of pH on the stability of plant phenolic compounds. *J Agric Food Chem* 48:2101-2110 (2000).
- 76 Masuda T, Mizuguchi S, Tanaka T, Iritani K, Takeda Y and Yonemori S, Isolation and structure determination of new antioxidative ferulic acid glucoside esters from the rhizome of *Alpinia speciosa*, a Zingiberaceae plant used in Okinawan food culture. *J Agric Food Chem* 48:1479-1484 (2000).
- 77 Kikuzaki H, Hisamoto M, Hirose K, Akiyama K and Taniguchi H, Antioxidant properties of ferulic acid and its related compounds. *J Agric Food Chem* 50:2161-2168 (2002).
- 78 Medina I, Tombo I, Satue-Gracia MT, German JB and Frankel EN, Effects of natural phenolic compounds on the antioxidant activity of lactoferrin in liposomes and oil-in-water emulsions. *J Agric Food Chem* 50:2392-2399 (2002).
- 79 Naim M, Striem BJ, Kanner J and Peleg H, Potential of ferulic acid as a precursor to off-flavours in stored orange juice. *J Food Sci* 53:500-512 (1988).
- 80 Narziss I, Miedaner H and Nitzsche F, Formation of 4-vinyl-guaiacol during production of Bavarian wheat beer. *Monatsschr Brauwiss* 43:96-100 (1990).
- 81 Lattanzio V, De Cicco V, Di Venere D, Lima G and Salerno M, Antifungal activity of phenolics against fungi commonly encountered during storage. *It J Food Sci* 1:23-30 (1994).
- 82 Ishihara M, Hasegawa M, Taira T and Toyama S, Isolation and antimicrobial activity of feruloyl oligosaccharide ester from pineapple stem residues. *J Jap Soc Food Sci Technol—Nippon Shokuhin Kagaku Kogaku Kaishi* 47:23-29 (2000).
- 83 Stead D, The effect of hydroxycinnamic acids and potassium sorbate on the growth of 11 strains of spoilage yeasts. *J Appl Bacteriol* 78:82-87 (1995).
- 84 Oosterveld A, Pol IE, Beldman G and Voragen AGJ, Isolation of feruloylated arabinans and rhamnogalacturonans from sugar beet pulp and their gel forming ability by oxidative cross-linking. *Carbohydr Polym* 44:9-17 (2001).
- 85 Micard V and Thibault JF, Oxidative gelation of sugar-beet pectins: use of laccases and hydration properties of the cross-linked pectins. *Carbohydr Polym* 39:265-273 (1999).
- 86 Oosterveld A, Beldman G and Voragen AGJ, Oxidative cross-linking of pectic polysaccharides from sugar beet pulp. *Carbohydr Res* 328:199-207 (2000).
- 87 Thibault JF, Garrpeau C and Durand D, Kinetics and mechanism of the reaction of ammonium persulfate with ferulic acid and sugar-beet pectins. *Carbohydr Res* 163:15-27 (1987).
- 88 Hurrell RF and Finot PA, Nutritional consequences of the reactions between proteins and oxidized polyphenolic acids. *Adv Exp Med Biol* 177:423-435 (1984).
- 89 Oudgenoeg G, Hilhorst R, Piersma SR, Boeriu CG, Gruppen H, Hessing M, Voragen AGJ and Laane C, Peroxidase-mediated cross-linking of a tyrosine-containing peptide with ferulic acid. *J Agric Food Chem* 49:2503-2510 (2001).
- 90 Kwok KC and Ou SY, Application of ferulic acid in the preparation of SPI-based edible film. *Sci Technol Food Ind* 129:24-27 (2002).
- 91 O'Connell JE and Fox PF, Proposed mechanism for the effect of polyphenols on the heat stability of milk. *Int Dairy J* 9:523-536 (1999).
- 92 Headley S and Massad S, *Nutritional supplements for athletes*, National Association for Sport and Physical Education, an Association of the American Alliance for Health, Physical Education, Recreation and Dance, pp 51-52 (1999).
- 93 Berning JR and Steen SN, *Nutrition for sport and exercise*, 2nd edn. Aspen Publishers Inc, Gaithersburg, Maryland, p 130 (1998).
- 94 Carbonare MD and Pathak MA, Skin photosensitizing agents and the role of reactive oxygen species in photoaging. *J Photochem Photobiol B* 14:105-124 (1992).
- 95 Podda M, Traber MG, Weber C, Yan LJ and Packer L, UV-radiation depletes antioxidants and causes oxidative damage in a model of human skin. *Free Radic Biol Medicine* 24:55-65 (1998).
- 96 RomeroGraillet C, Aberdam E, Clement M, Ortonne JP and Ballotti R, Nitric oxide produced by ultraviolet-irradiated keratinocytes stimulates melanogenesis. *J Clin Invest* 99:635-642 (1997).
- 97 Saija A, Tomaino A, Trombetta D, De Pasquale A, Uccella N, Barbuzzi T, Paolino D and Bonina F, *In vitro* and *in vivo* evaluation of caffeic and ferulic acids as topical photoprotective agents. *Int J Pharm* 199:39-47 (2000).
- 98 Murakami A, Nakamura Y, Koshimizu K, Takahashi D, Matsumoto K, Hagihara K, Taniguchi H, Nomura E, Hosoda A, Tsuno T, Maruta Y, Kim HW, Kawabata K and Ohigashi H, FA15, a hydrophobic derivative of ferulic acid, suppresses inflammatory responses and skin tumor promotion: comparison with ferulic acid. *Cancer Lett* 180:121-129 (2002).